Journal of *Fungi*

MDPI

Article

Comparative Genomics Reveals Host-Specific Adaptation of *Pyricularia oryzae* Strains Isolated from Rice and Barnyard Grass

Wenda Sun [1,†], Xiaohan Zhang [1,†], Zhuan Zhang [1], Xiaofang Xie [1], Song Tang [1], Tian Song [1], Baoxu Lu [1], Jiafeng Wang [2], Zhibin Liang [1], Xiaofan Zhou [1,*] and Yizhen Deng [1,*]

[1] Guangdong Provincial Key Laboratory of Microbial Signals and Disease Control, Integrative Microbiology Research Centre, South China Agricultural University, Guangzhou 510642, China
[2] Nation Engineering Research Center of Plant Space Breeding, South China Agricultural University, Guangzhou 510642, China
* Correspondence: xiaofan_zhou@scau.edu.cn (X.Z.); dengyz@scau.edu.cn (Y.D.)
† These authors contributed equally to this work.

Abstract

Barnyard grass, a widespread and persistent weed in rice paddies, belongs to the same family as rice and may act as a bridge host for the rice blast fungus. This study utilized comparative genomics to analyze six *Pyricularia oryzae* strains isolated from barnyard grass (Baicao series) and rice (GDYJ7 and ZJX18), integrating pathogenicity assays, whole-genome sequencing, and functional annotation. Pathogenicity tests demonstrated host specificity, as Baicao series strains caused typical lesion symptoms on barnyard grass but not on rice leaves, while GDYJ7 and ZJX18 caused lesions mainly on rice. Genomic analyses indicated that Baicao series strains possessed larger genomes (41.04 Mb to 41.16 Mb) with a higher content of repetitive sequences (6.68% to 7.09%) compared to rice strains GDYJ7 and ZJX18 (38.69 Mb and 39.05 Mb; 3.66% and 3.71% repeats). Phylogenetic analysis confirmed that Baicao series strains represent a grass-infecting pathotype of *P. oryzae* species, as they were grouped with the established grass-isolated *P. oryzae* strains, while GDYJ7 and ZJX18 were grouped with rice-isolated *P. oryzae* strains. However, Baicao series, GDYJ7 and ZJX18 are all relatively distant from *P. grisea* species. PCR amplification revealed that Baicao series strains harbored significantly fewer avirulence genes (*Avr-Pib*, *Avr-Pizt*, *PWL3*) than GDYJ7 and ZJX18 (*Avr-Pib*, *Avr-Pizt*, *Avr-Pi9*, *Avr-Pik*, *PWL2*), with Baicao9 retaining only *Avr-Pib*. In summary, our results suggested that the genomic sequences of the barnyard grass-isolated strains serve as a valuable resource for the study of *P. oryzae* strains with differential host preference and provide novel insights into the evolution of pathogen genomes during host adaptation.

Keywords: *Pyricularia*; whole-genome sequencing; avirulence genes; comparative genomics

Check for updates

Academic Editor: Willem JG Melchers

Received: 11 January 2026
Revised: 31 January 2026
Accepted: 2 February 2026
Published: 5 February 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Rice (*Oryza sativa* L.) is one of the most important food crops for people worldwide [1]. Rice blast caused by *Pyricularia oryzae* (syn. *Magnaporthe oryzae*) is one of the most devastating diseases affecting rice crops worldwide, leading to 10% to 30% losses of global rice yield every year [2–4]. In the past three decades, rice blast has been developed as a model system to study fungal–plant interactions [5,6]. *P. oryzae* was the first plant pathogenic fungus to have its genome sequenced and made available to the public [7]. *Pyricularia* species can infect not only cereal crops such as rice, wheat [8], and barley [9] but also a wide range of grasses including crabgrass (*Digitaria* spp.), foxtail (*Setaria* spp.), and

ryegrass (*Lolium* spp.) [10–12]. Studies have also shown that certain *Pyricularia* species are capable of infecting non-gramineous hosts such as banana (*Musa* spp.) and ginger (*Zingiber* spp.) [13,14]. Host shift and host expansion by fungal plant pathogens have increased the frequency of emerging pathogens and the incidence of disease across crops, posing a serious threat to global food security [15–17]. A notable case of host shift and co-evolution of this fungus in recent decades is the outbreak of wheat blast caused by *P. oryzae* in South America in the early 1990s [18]. Additionally, the infection of wheat plants displaying white head symptoms in Kentucky, USA, has been suggested as another key example of host shift, where the pathogen likely jumped from ryegrass (*Lolium* spp.) to wheat as its new host [19]. Another study suggests that *P. oryzae* may shift from millet (*Setaria* spp.) to rice [20]. The broad host range of *P. oryzae*, coupled with its potential for host shift, significantly contributes to the emergence and outbreak of novel diseases and is thus worthy of in-depth investigation on the molecular mechanism of host adaptation and expansion of the *P. oryzae* population. Availability of genome sequences of *P. oryzae* population thus provides the basis for making such analyses.

The publication of the *P. oryzae* strain 70-15 genome has since spurred the sequencing and assembly of numerous field isolates [7], such as Guy11 [21], B157 [22], Y34 [23], and 98-06 [24]. Comparative genomic and functional studies of these strains have uncovered substantial genome plasticity and highlighted the contribution of lineage-specific genes to pathogenicity [25–27]. More recently, advances in sequencing technologies have enabled large-scale sequencing of field isolates from rice as well as isolates collected from various grass and cereal hosts [28,29]. Population-level analyses of these genomes indicate that rapidly evolving genes acts as a key driver of pathogen specialization [30–32]. Nevertheless, the genomic landscape of most species within the *Pyricularia* spp. remains largely unexplored. At present, genomic information on rice-infecting *Pyricularia* strains is abundantly available but less reported in the Poaceae grass-infecting *Pyricularia* strains. Therefore, comparative genomic analysis of *P. oryzae*, the fungal pathogen responsible for rice blast, could aid in monitoring its host shift and/or spread, thereby helping to predict and prevent potential epidemics.

Pathogenic strains of *Pyricularia* contain more genes encoding secretory proteins and avirulence (*Avr*) genes in their genomes compared to non-pathogenic strains, which may contribute to their successful host invasion [33]. The interaction between rice and the rice blast fungus follows the classic gene-to-gene interaction theory [34]. The host specificity of the plant pathogen is determined by the interaction between the plant host's resistance (*R*) genes and the fungal pathogen's *Avr* genes [35]. To date, more than 40 *Avr* genes have been identified in the rice blast fungus, 12 of which have been cloned [36]. Variation in *Avr* genes leads to differences in virulence among strains of the fungus [37]. Mechanisms such as gene deletion, frameshift mutations, and transposon insertion can alter the function of *Avr* genes, thereby affecting the pathogenicity of strains carrying such *Avr* genes [38,39]. For example, Pot3 transposon has been identified as a major reason to generate different variations of *Avr-Pizt* and *Avr-Pib*. The insertion of Pot3 transposon at the −462 bp site in the promoter region of the *AvrPiz-t* gene enabled the *P. oryzae* strains Guy11 to evade recognition by rice carrying the *Piz-t* resistance gene, thereby causing a shift from an "avirulent" to a "virulent" phenotype [40]. In all 248 *P. oryzae* isolates from Philippines, insertions of the Pot3 transposon were identified within the *AvrPib* gene, mainly occurring at three distinct sites: two in the promoter region (at positions −304 bp and −125 bp) and one within the coding region (at 169 bp). Such Pot3 insertion resulted in the failure of *AvrPib* gene expression, and consequently, to the evasion of recognition by the rice *Pib* resistant gene, leading to a shift from an "avirulent" to a "virulent" phenotype [41]. This type of genetic variation provided the *P. oryzae* population with a rapid adaptive advantage, thereby enabling it to overcome

crop resistance. Therefore, monitoring the loss and variation of *Avr* genes in populations of the rice blast fungus is crucial for safeguarding rice production in the region.

Barnyard grass, a weed commonly found in and around rice fields, belongs to the same family as rice (Poaceae). It may serve as a bridge host for the rice blast fungus, facilitating the survival and dispersal of the pathogen. In this study, four *P. oryzae* strains were isolated from blast-like lesions on barnyard grass growing near paddy fields at the Baisha Experimental Base (111°50′ E, 21°52′ N) in Yangjiang, Guangdong Province, China. Together with two previously isolated rice-infecting *P. oryzae* strains, GDYJ7 and ZJX18, from *Xian* rice (Indica) in Guangdong and Zhejiang province, respectively, we assessed the pathogenicity of these six strains on detached leaves of barnyard grass and rice and concurrently examined the presence or absence of several known *Avr* genes. To further investigate their taxonomic classification and pathogenic mechanisms at the genomic level, we performed whole-genome sequencing on all six isolates. Overall, this study aims to reveal the molecular mechanism of host adaptation by different strains within the *P. oryzae* population.

2. Results

2.1. Isolation and Identification of Six Pyricularia Strains from Barnyard Grass and Rice with Leaf Blast Symptoms

We isolated a total of four *Pyricularia* single-spore strains from the same diseased barnyard grass leaves collected near rice fields in Yangjiang, designating them Baicao4, Baicao5, Baicao6, and Baicao9. In addition, we included the previously isolated *P. oryzae* strains GDYJ7 from diseased rice in Yangjiang and ZJX18 from diseased rice in Zhejiang Province (Figure 1A,B). On RPA medium, strains of the Baicao series formed white, fluffy colonies with dense mycelium. Strain GDYJ7 developed grayish-black, fluffy colonies with moderately dense mycelium, while strain ZJX18 produced grayish-white colonies that developed black speckles at later stages, accompanied by relatively sparse mycelium (Figure 1C). The conidia of all the above-mentioned strains were pyriform in shape and contained two septa (Figure 1C).



Figure 1. Isolation and morphological characteristics of six *Pyricularia* strains. (**A**) Lesion symptoms on barnyard grass or rice leaves in the field. (**B**) Single-spore isolation workflow. (**C**) Colony and conidia morphology. Scale bars = 10 μm. All strains were grown for 7 days on RPA medium before being photographed.

2.2. Pathogenicity Analysis of the Isolated Strains

We conducted pathogenicity assays on these six strains, among which the pathogenicity of GDYJ7 was validated [42,43] and therefore served as a positive control for rice blast. The results (Figure 2A,B) showed that all Baicao strains successfully infected barnyard grass (with Baicao9 showing weaker pathogenicity) and caused typical lesion symptom but showed almost no infection on rice. Strain GDYJ7 was able to infect the susceptible rice cultivars CO39 and NPB, exhibited limited infection on the moderately resistant cultivar ZH11, and failed to infect barnyard grass. Strain ZJX18 was the most virulent, capable of infecting all three rice cultivars (CO39, NPB, and ZH11) with prominent lesions, and also caused mild lesions on barnyard grass (Figure 2A,B). These findings confirm that isolates from different host origins exhibit clear host preference.



Figure 2. Pathogenicity assay of the strains. (**A**) Spore suspensions (1×10^5/mL) were inoculated on the detached leaves of barnyard grass and different rice varieties (ZH11, NPB, and CO39). *P. oryzae* strain GDYJ7 served as a positive control and H_2O as a blank control. Scale bars = 1 cm. (**B**) Quantitative analysis of lesion area based on the results of (**A**). The mean ± S.D. was derived from three independent biological repeats. Statistical analysis was performed using a two-tailed unpaired Student's *t*-test versus GDYJ7. ns, **, ***: no significance, $p < 0.01$, $p < 0.001$, respectively.

2.3. Genome Sequencing and Assembly

Next, we performed whole-genome sequencing for the six *Pyricularia* strains using Illumina technology. After quality control of the raw data, de novo assembly generated genome sequences of Baicao4 (size: 41.04 Mb, 1339 contigs), Baicao5 (41.18 Mb, 1278 contigs), Baicao6 (41.44 Mb, 1149 contigs), Baicao9 (41.46 Mb, 1117 contigs), GDYJ7 (38.69 Mb, 1358 contigs), and ZJX18 (39.05 Mb, 1463 contigs). The Contig N50 values ranged from 99,612 bp to 141,887 bp across the six strains, indicating reasonable levels of genome assembly continuity. The GC contents of the four Baicao series strains (50.15% to 50.31%)

were slightly lower than those of the two rice-isolated strains GDYJ7 and ZJX18 (51.24% and 51.22%) (Table 1).

Table 1. Genome assembly statistics.

Features	Baicao4	Baicao5	Baicao6	Baicao9	GDYJ7	ZJX18
All Length (Mb)	41.04	41.18	41.44	41.46	38.69	39.05
Contig num	1339	1278	1149	1117	1358	1463
Contig max len (bp)	732,390	477,166	581,875	627,432	731,285	688,677
Contig average len (bp)	30,650.10	32,219	36,061.90	37,118.60	28,487.50	26,690.40
Contig N50 (bp)	99,612	92,052	119,374	102,581	147,078	141,887
GC Ratio	50.31%	50.26%	50.16%	50.15%	51.24%	51.22%

Assessment with BUSCO showed that all strains contained at least 4309 (95.9%) complete single-copy genes, indicating high levels of genome completeness (Table 2). Gene prediction analysis identified 11,114 to 11,440 protein-coding genes in the genomes of the six strains (Table 3). Repetitive sequences are widespread in eukaryotic genomes, and their proportion is closely associated with genome size [44]. In this study, repetitive sequence annotation revealed higher levels of repeat content in the four Baicao strains (6.68% to 7.09%) relative to GDYJ7 and ZJX18 (3.66% and 3.71%) (Table 3). Transposable elements represent a major component of repetitive sequences and can drive gene evolution through chromosomal rearrangements and other forms of genomic variation [45]. A total of 650 to 760 DNA transposons were predicted in all six strains genomes, accounting for less than 1% of the total length of each genome. Notably, long interspersed nuclear elements (LINEs) were present in all six strains, whereas short interspersed nuclear elements (SINEs) were only detected in Baicao4, Baicao5, GDYJ7, and ZJX18. In addition, long terminal repeat (LTR) retrotransposons of the Gypsy/DIRS1 and Ty1/Copia types were identified in all six strains, while the BEL/Pao type of LTR retrotransposons was found in strain Baicao9 only (Table 3).

Table 2. Genome assembly evaluation.

Features	Baicao4	Baicao5	Baicao6	Baicao9	GDYJ7	ZJX18
Complete and single-copy	4315 (96.1%)	4316 (98.0%)	4319 (98.1%)	4319 (97.8%)	4321 (96.2%)	4309 (95.9%)
Complete and duplicated	4 (0.1%)	4 (0.1%)	4 (0.1%)	4 (0.1%)	4 (0.1%)	5 (0.1%)
Fragmented	90 (2.0%)	92 (2.0%)	87 (1.9%)	89 (0.7%)	87 (1.9%)	81 (2.0%)
Missing (not recovered in assembly)	83 (1.8%)	80 (1.8%)	82 (1.8%)	83 (1.2%)	80 (1.8%)	87 (1.9%)
Total BUSCO	4492 (100%)	4492 (100%)	4492 (100%)	4492 (100%)	4492 (100%)	4492 (100%)

Non-coding RNAs (ncRNAs) are key regulators involved in cellular processes including growth, differentiation, survival, and apoptosis [46]. Our prediction across the six genomes identified all major ncRNA types, including transfer RNAs (tRNAs; 215 to 320), ribosomal RNAs (rRNAs; 40 to 54), small nuclear RNAs (snRNAs, 22 in each strain), and small RNAs (sRNAs, four in each strain) (Table 3).

Table 3. Genome component analysis.

Prediction Features	Baicao4	Baicao5	Baicao6	Baicao9	GDYJ7	ZJX18
Gene Number	11,114	11,116	11,151	11,171	11,440	11,118
tRNA	215	216	216	216	312	320
rRNA	53	52	40	47	48	54
sRNA	4	4	4	4	4	4
snRNA	22	22	22	22	22	22
DNA transposons	691	697	730	729	650	760
LINE	121	142	146	112	380	404
SINE	8	8	-	-	9	9
LTR	1954	2250	1732	1890	1370	1347
BEL/Pao	-	-	-	8	-	-
Gypsy/DIRS1	1210	1573	707	1255	1192	1181
Ty1/Copia	457	455	972	444	178	166
Effect Protein Number	3176	3170	3189	3191	3097	3172
Cytoplasmic effector	2763	2751	2770	2776	2715	2783
Apoplastic effector	413	419	419	415	382	763

"-" stands for non-existence.

2.4. Phylogenetic Analyses

Comparative genomic analysis based on average nucleotide identity (ANI) revealed that the four Baicao series strains shared 99.05% to 99.07% identity with *P. oryzae* strain D15/s47. Strain GDYJ7 showed 99.74% and 99.69% identity to *P. oryzae* strain 70-15 and P131, respectively, and strain ZJX18 with 99.75% and 99.82% identity with 70-15 and P131 (Table S1). We previously reported a barnyard grass isolated *P. grisea* strain SCAU-2 [29]. Here, we found that Baicao series strains shared only 86.74 to 86.75% and 86.90 to 86.94% identity with two *P. grisea* strain SCAU-2 and NI907, respectively (Table S1).

We further constructed a maximum-likelihood (ML) phylogenetic tree using single-copy orthologous gene sequences. The Baicao series strains clustered within the same evolutionary clade as *P. oryzae* E34, MZ5-1-6 (isolated from *Eleucine*), and *P. grisea* D15 (isolated from *Digitaria*). The strains GDYJ7 and ZJX18 grouped together with the *P. oryzae* strains 70-15, Y34, and ZJG1 (isolated from rice). SCAU-2 was grouped with *P. grisea* strains NI907 and D1, separated from the clade containing Baicao series strains (Figure 3). Interestingly, the Baicao series strains are closer to the *P. oryzae* strains preferring the host Eleusine but relatively distant from the *P. grisea* strains preferring Digitaria. Therefore, we conclude that Baicao series strains belong to the grass pathotype of *P. oryzae* species, rather than *P. grisea* species.

2.5. Prediction and Analyses of Pathogenicity-Related Genes

To identify potential virulence genes in the fungal pathogens, we performed gene annotation using the Pathogen–Host Interactions database (PHI) and the Database of Fungal Virulence Factors (DFVF). Based on the PHI database, 1416, 1430, 1417, 1412, 1422, and 1434 PHI-related genes were annotated in strains Baicao4, Baicao5, Baicao6, Baicao9, GDYJ7, and ZJX18, respectively. Among these, the numbers of genes associated with increased pathogenicity (enhanced virulence) were 19, 19, 19, 19, 18, and 17, respectively (Figure 4A). Details of PHI-related genes with sequence identity > 90% are provided in Table S2. According to the DFVF database, 1258 (11.3%), 1290 (11.6%), 1294 (11.6%), 1287 (11.2%), 1288 (11.6%), and 1293 (11.6%) protein-coding genes were successfully annotated in strains Baicao4, Baicao5, Baicao6, Baicao9, GDYJ7, and ZJX18, respectively (Figure 4B). Detailed information on DFVF-related genes with sequence identity > 90% is listed in Table S3.



Figure 3. Phylogenetic analysis. Constructing a phylogenetic tree based on single-copy orthologous gene sequences using the maximum-likelihood (ML) method, and the ANI analysis was integrated for validation. The red typeface represents the strains isolated in this study. A heatmap of the ANI values between *P. oryzae* 70-15/*P. pennisetigena* Br36/*P. grisea* NI907 and the 17 representative *Pyricularia* strains is shown on the right side of the tree.



Figure 4. Comparative analysis of gene annotations and plant cell wall-degrading enzymes for six strains based on the PHI, DFVF, CAZy, and PCWDEs databases. (**A**) Annotation statistics of the six pathogenic strains in the PHI database. (**B**) Proportion of genes annotated in the DFVF database for the six strains. (**C**) CAZy functional annotation results for the six strains. (**D**) Comparative analysis of fungal PCWDEs among the six strains.

Additionally, carbohydrate-active enzymes are closely linked to fungal growth, development, and pathogenicity [47,48]. Based on the Carbohydrate-Active Enzymes (CAZy) database, strains Baicao4, Baicao5, Baicao6, Baicao9, GDYJ7, and ZJX18 were annotated with 2392, 2399, 2399, 2405, 2411, and 2439 related genes, respectively. The CAZyme annotation profiles of the six strains were largely similar in composition and quantity. Among these, glycoside hydrolases (GHs) were the most abundant (1160 to 1183), which was followed by glycosyltransferases (GTs; 548 to 557). Polysaccharide lyases (PLs) represented the lowest proportion among all enzyme classes (Figure 4C). The functional composition of plant cell wall-degrading enzymes (PCWDEs) across the six strains exhibited overall high consistency (Table S4), with hemicellulases predominating, followed by cellulases and lignin-related enzymes. Only in strain GDYJ7 was there a slight reduction in the proportion of hemicellulases (Figure 4D).

2.6. Prediction and Analyses of Membrane Transport Proteins

Membrane transport proteins play a crucial role in pathogen–host interactions. In this study, we predicted 898, 900, 905, 898, 903, and 911 membrane transport proteins in strains Baicao4, Baicao5, Baicao6, Baicao9, GDYJ7, and ZJX18, respectively (Table 4). Among these membrane transport proteins, the composition was generally similar across the six strains. In the four Baicao series strains, secondary transporters represented the highest proportion, with counts of 232, 236, 236, and 228, respectively. In strains GDYJ7 and ZJX18, the numbers of secondary transporters were 228 and 233, respectively. Next, primary active transporters showed a proportion largely comparable to that of secondary transporters. The six strains contained 226, 226, 226, 226, 231, and 231 primary active transporters, respectively (Figure 5A).

Table 4. The number of transporters.

Transporters	Baicao4	Baicao5	Baicao6	Baicao9	GDYJ7	ZJX18
ABC	43	43	42	42	45	42
MFS	66	67	70	60	64	69
other	789	790	793	788	794	800
total	898	900	905	890	903	911

The number of major facilitator superfamily (MFS) transporters outnumbered that of ATP-binding cassette (ABC) transporters in all six strains: the counts of MFS transporters in Baicao4, Baicao5, Baicao6, Baicao9, GDYJ7, and ZJX18 were 66, 67, 70, 60, 64, and 69, respectively, whereas the corresponding numbers of ABC transporters were 43, 43, 42, 42, 45, and 42 (Table 4).

Both the ABC and MFS superfamilies contain multidrug transporters, which help pathogens resist adverse environmental stresses [49]. Among the six strains in this study, multidrug transporters accounted for the largest proportion and showed relatively similar abundance. In terms of specific subtypes, the most abundant multidrug transporters in all six strains were proteins of the DHA1 family and multidrug resistance (MDR) proteins, followed by DHA2 family proteins and pleiotropic drug resistance (PDR) proteins (Figure 5B).



Figure 5. Prediction of membrane transport proteins and secondary metabolite synthases. (**A**) Annotation statistics of the six pathogenic strains in the TCDB database. (**B**) Comparative analysis of multidrug transporters from the ABC and MFS families in the six pathogenic strains. (**C**) Statistical results of secondary metabolite biosynthetic gene clusters in the six pathogenic strains.

2.7. Prediction and Analyses of Secondary Metabolite Biosynthetic Gene Clusters

Plant pathogens produce a wide array of secondary metabolites, which can serve as virulence factors and facilitate host infection [50,51]. In this study, we identified 71, 70, 70, 71, 74, and 83 secondary metabolite biosynthetic gene clusters in the genomes of strains Baicao4, Baicao5, Baicao6, Baicao9, GDYJ7, and ZJX18, respectively. The composition of these clusters was largely similar across strains, with terpene and T1PKS clusters being the most abundant; each strain contained 18–21 clusters of these two types. NRPS and NRPS-like clusters varied somewhat among strains: GDYJ7 and ZJX18 possessed relatively fewer NRPS clusters, while NRPS-like clusters were slightly more numerous in GDYJ7. In contrast, indole clusters were low in number and relatively consistent in all strains (Figure 5C).

2.8. Presence and Variation of Avr Genes

We successfully obtained the sequences of six established *Avr* genes, *Avr-Pib*, *Avr-Pizt*, *Avr-Pi9*, *Avr-Pik*, *PWL2*, and *PWL3*, from the whole genomes of the six strains. We further verified these *Avr* genes by PCR amplification, using the primers listed in Table 5. Combined with genome sequencing and PCR amplification, we found that Baicao4, Baicao5, and Baicao6 shared a conserved set of *Avr* genes including *Avr-Pib*, *Avr-Pizt*, and *PWL3* but lacked *Avr-Pik* and *PWL2*. However, Baicao9 strain contained only *Avr-Pib*, exhibiting marked uniqueness compared to the other Baicao strains. This suggests a potential divergence in pathogenic mechanisms or host adaptation compared to the other barnyard grass isolates. In stark contrast, the two rice-isolated strains GDYJ7 and ZJX18 possessed *Avr-Pib*, *Avr-Pizt*, *Avr-Pik*, *Avr-Pi9*, and *PWL2*. *PWL3* was missing in GDYJ7 but present in ZJX18 (Figure S1; Table 6). Two Avr genes, *Avr-Pia* and *Avr-Pita2*, were detected in our

previously isolated *P. grisea* strain SCAU-2 [29] but not found in genomes of the six strains in this study, and neither did they amplified by PCR (Figure S1).

Table 5. Primers used in this study.

Gene Name	Primer Sequence (5′-3′)	Source
Avr-Pia	F: CATCGCTTTGCCCTCATT R: ACTTGATTCCTCCCGTAAACAG	This study
Avr-Pib	F: AAGTCCTTCCCATTACCCTA R: GCAATAACCATCCAGCCATA	[52]
Avr-Pizt	F: GATCAAATGAACACCAGGAA R: CGATGAAGAATGGAAGAATG	[53]
Avr-Pi9	F: CCTTCTAGTCATTCCTTTGG R: AGGCGAATGTGCTTACTACT	This study
Avr-Pik	F: AATTTATTCAACTGCCACTCTG R: AACCTCGTCAAACCTCCCTA	[52]
Avr-Pita2	F: TTTCGGCCCAACTCCGGTCC R: TAAAGGGTCCACTGACCCCG	[54]
PWL2	F: ATGAAATGCAACAACATC R: CCTCACACTTAAGTTAACAC	[55]
PWL3	F: GCGTGCTCATTTGTAAACC R: TTCCTTCATTTCTCTCCCTG	This study

Table 6. Avirulence genes distribution.

Strains	*Avr-Pib*	*Avr-Pizt*	*Avr-Pi9*	*Avr-Pik*	*PWL2*	*PWL3*
Baicao4	+	+	-	-	-	+
Baicao5	+	+	-	-	-	+
Baicao6	+	+	-	-	-	+
Baicao9	+	-	-	-	-	-
GDYJ7	+	+	+	+	+	-
ZJX18	+	+	+	+	+	+/aa sequence altered; remature termination

"-" stands for non-existence. "+" stands for existence.

We aligned the sequences of *Avr-Pib*, *Avr-Pizt*, and *PWL3* shared by the isolates from barnyard grass or rice obtained in this study, the *P. oryzae* strains from different hosts (*Oryza*, *Triticum*, *Eleusine*), and *P. grisea* strains infecting *Digitaria* (if available). Simultaneous alterations at two sites, A137T and A158C, were found in the coding sequences of *Avr-Pib* in the four Baicao series strains and their closest strain MZ5-1-6 (based on phylogenetic analysis showed in Figure 3), resulting in amino acid changes as E46Vand Y53S, respectively, compared to other strains (Figure S2). Two *P. grisea* strains displayed multiple point mutations that are distinct from *P. oryzae* strains, but at A103G, G178A, and A213T they are same as one of the wheat-infecting strain 131021 (Figure S2). *Avr-Pizt* is absent in *P. griase* strains and Baicao9. Altered sites in the *Avr-Pizt* coding sequences could clearly distinguish the *P. oryzae* strains infecting rice, wheat, and grass (Figure S3). The coding sequences of the *PWL3* gene are present in three Baicao strains (except for Baicao9) and several other *P. oryzae* strains. Interestingly, we noticed a clear pattern of nucleotide point mutations, and corresponding changes in amino acid sequences, that could divide all the tested strains into three groups: Baicao series strains and MZ5-1-6 in a group (group A) that is closer with four rice-infecting strains (group B), while the two wheat-infecting strains are grouped with the other grass-infecting strains (group C) (Figure S4). Based on these observations,

we hypothesized that the host range of *P. oryzae* populations could be inferred based on a combination of *Avr* gene variations.

In summary, differences in the *Avr* gene profiles between the barnyard grass-isolated Baicao strains and the rice-isolated strains (GDYJ7, ZJX18), as well as established *P. grisea* strain (SCAU-2), may reflect differential adaptation strategy towards different hosts.

3. Discussion

This study conducted an integrated comparative genomic analysis of six *P. oryzae* strains obtained from barnyard grass and rice. It revealed dynamic changes and specific adaptive strategies at the genomic level during the pathogen's adaptation to different Poaceae hosts. Our findings not only confirmed the central role of host specificity in the evolution of the blast fungus but also provided novel genomic perspectives for understanding host jump and potential epidemic risks associated with this important pathogen.

First, we confirmed a clear correlation between the host origin and pathogenic phenotype. The Baicao series strains from barnyard grass successfully infected their original host but were nearly non-pathogenic to rice. In contrast, the rice-isolated strains GDYJ7 and ZJX18 exhibited strong pathogenicity towards rice but showed no or mild infection capability on barnyard grass (Figure 2). Phylogenetic analysis grouped the Baicao series strains with *P. oryzae* strains isolated from Eleusine, while GDYJ7 and ZJX18 clustered with typical rice-infecting *P. oryzae* strains (Figure 3), all separated from the *P. grisea* strains NI907 and SCAU-2. Eleusine genus is represented by species *Eleusine indica* and *Eleusine coracana*. *E. indica* is a widely distributed weed in many regions [56], while *E. coracana* is utilized as a dual-purpose grain and forage plant in Africa [57]. Our Baicao series strains cluster phylogenetically with multiple *E. coracana*-isolated strains, suggesting that, during evolution, some *P. oryzae* strains may have utilized grasses as hosts before they gained ability to infect grain crops. This supported their classification into different pathotypes from an evolutionary relationship perspective. Barnyard grass, a pervasive and difficult-to-eradicate weed in paddy ecosystems, has long been considered a potential alternative host and "refuge" for *P. oryzae* [10,58]. Previous studies have indicated the existence of gene flow between pathogens on grasses and cereals [59], and host jump is a key factor in the outbreak of emerging diseases such as wheat blast [19,60]. In this study, interestingly, the rice-infecting *P. oryzae* strain ZJX18 caused a mild lesion symptom on grass (Figure 3). This could be either a residual infectivity to an old host after it jumped from grass to rice, or alternatively, it may broaden its host range from rice to the grass growing at the rice paddy.

Second, analysis of genomic components revealed potential genomic plasticity associated with host adaptation. Transposable elements (TEs), as major drivers of repetitive sequences and genomic plasticity, likely played a key role through differences in their abundance and type [61,62]. For instance, BEL/Pao-type LTR retrotransposons were detected exclusively in the strains Baicao9. Existing research suggests that host specialization is linked to TE-associated dynamic gene gain and loss [26]. In this study, the genome size of barnyard grass-isolated strains was slightly larger than that of rice-isolated strains (Table 1), and their repetitive sequence proportion was significantly higher than that of rice-isolated strains (Table 3). Active TEs might have rapidly promoted genetic variation within pathogen populations by causing chromosomal rearrangements, gene inactivation, or novel gene generation, thereby facilitating adaptation to different host selection pressures [63,64]. For example, Pot3 has been identified as a major reason to generate different variations of *Avr-Pizt* and *Avr-Pib* [40,41]. Studies on other plant pathogens, such as *Fusarium* species causing Fusarium head blight (FHB) or rust fungi causing wheat stem rust, have also shown that rapid changes in transposon-rich genomic regions constitute a common strategy for pathogen genome evolution and rapid adaptation to different environments [65–67].

Furthermore, annotation of pathogenesis-related genes indicated that despite host specialization, strains from different origins maintained a high degree of conservation in core pathogenic mechanisms. Annotations based on the PHI and DFVF databases showed that all strains possessed a similar number of pathogenesis-related genes and virulence factors. The composition and quantity of CAZymes were also highly similar among strains, with GHs being the most abundant family (Figure 4). This implied that fundamental infection strategies, such as plant cell wall degradation, represented a common toolkit for *Pyricularia* pathogens infecting different Poaceae hosts. However, finer regulation might have occurred at the level of effector secretion and secondary metabolite synthesis [68,69]. Membrane transport proteins, particularly multidrug transporters from the ABC and MFS families, are crucial in pathogen–host interactions. They might have been involved in effector secretion or helped the pathogen counteract antimicrobial substances produced by the host [28,70,71]. For instance, *Botrytis cinerea* AI18 neutralizes phytoalexins from Solanaceae and Fabaceae plants via a dual mechanism involving the efflux transporter BcatrB and complementary metabolizing enzymes [72]. Despite no significant difference in the number of transporters between the two types of *Pyricularia* strains, differences in their expression levels might determine their differential roles in effector secretion or host infection. The accumulation of diterpenoid phytoalexins (DPs) in rice, notably momilactones and phytocassanes, is induced upon infection by the blast fungus [73]. Therefore, *P. oryzae* may enhance its survival during infection by upregulating the expression of multidrug transporters, such as those from the ABC and MFS families, to efflux toxic compounds like DPs out of the cell. Fungi are capable of producing a vast array of secondary metabolites, which serve various functions, such as acting as signaling molecules for inter-microbial communication or as virulence factors in interactions with plant and animal hosts [74]. In this context, differences in secondary metabolite biosynthesis gene clusters, such as the relatively lower number of NRPS clusters in GDYJ7 and ZJX18, might have indicated that different strains synthesized distinct specific toxins or signaling molecules to adapt to their respective host environments (Figure 5C). The Baicao series strains derived from barnyard grass may harbor more diverse NRPS clusters, enabling the synthesis of compounds specifically adapted to the paddy field companion weeds host. This genetic capacity could enhance their colonization or virulence on this specific host.

The rice-isolated strains GDYJ7 and ZJX18 harbored a more diverse repertoire of *Avr* genes, including *Avr-Pi9*, *Avr-Pizt*, *Avr-Pik*, and *PWL2*. In contrast, less *Avr* genes were detected in the Baicao series strains, among which Baicao9 retains only *Avr-Pib* (Table 5). Products of the *Avr* genes were initially acting as a virulence factor towards the host lacking the corresponding *R* genes. Therefore, variations of the *Avr* gene coding sequences may be attributed to host adaptation and provide clues for origin of wheat- or rice-infecting strains from the *P. oryzae* population. *Avr-Pizt* is a small secreted peptide consisting of 108 amino acids [40], targeting the rice E3 ubiquitin ligases APIP6 and APIP10 to modulate plant defense responses and playing a crucial role in virulence [75–77]. *Avr-Pik* is a 113-amino acid protein with a 21-amino acid signal peptide and exhibits high variability, with multiple allelic forms present [78]. In susceptible rice (without corresponding R gene), *Avr-PikD* (the ancestral allele of *Avr-Pik*) inhibits the activity of the rice LSD1 transcriptional activator AKIP30, thereby facilitating *P. oryzae* infection via effector-triggered susceptibility (ETS) [79]. Both *Avr-Pib* and *Avr-Pi9* exhibit multiple types of variations, such as TEs insertions, point mutations, or deletions, which constitute the primary mechanisms leading to the loss of their avirulence function [36]. *Avr-Pib* weakens rice blast resistance by inhibiting the interaction between OsMAPKKK72 and OsMKK9, thereby reducing MAPK activation [80]. *Avr-Pi9* may function in compromising the basal resistance in rice by targeting the rice RING-type E3 ubiquitin ligase OsRGLG5 [81]. *PWL2* and *PWL3* are members of a

dynamic, rapidly evolving gene family, functioning as host-specificity determinants [82,83]. The effector protein *PWL2* acts as a virulence factor that suppresses plant immunity by interfering with the localization of the host protein HIPP43 at plasmodesmata [84]. In this study, we observed clear patterns in *Avr-Pizt* and *PWL3* coding sequence variation reflecting host preference to rice, wheat, or grasses (Figures S3 and S4). Furthermore, for *PWL3* coding sequences, Baicao series strains and MZ5-1-6 are closer with rice-infecting strains, while other grass-infecting strains were closer to wheat-infecting strains (Figure S4). The *PWL3* gene (absent in Baicao9) in the Baicao series strains and MZ5-1-6 was identical to the reference sequence U36995.1 that was reported from grass-isolated *P. grisea* strain, indicating that it may play a role in pathogenicity towards grass. Further mutations in the *PWL3* gene differentiate rice-infecting strains (including ZJX18) from Baicao series strains and MZ5-1-6 (Figure S4). Another rice-infecting strain, GDYJ7, lacks the *PWL3* gene. We infer that the point mutations of the *PWL3* gene, leading to alteration of amino acid sequence and premature termination in rice-infecting strains (Figure S4; Table 6), may be responsible for their loss of pathogenicity towards grass. To support this hypothesis, the ZJX18 strain showed residual infectivity to grass while GDYJ7 was completely non-pathogenic to grass; absence of *PWL3* in Baicao9 may be responsible for its weak pathogenicity to grass, compared to other Baicao strains (Figure 2). The presence of *Avr-Pi9* and *Avr-Pik* in GDYJ7 and ZJX18, but absence in Baicao series strains, may be a reason for higher virulence of the rice-isolated strains than Baicao series strains towards the rice host (without the corresponding *R* genes). In addition, presence of *PWL2* in GDYJ7 and ZJX18 may be a reason for their incapability to infect grass hosts, as reported [70]. The presence of *PWL3* in Baicao series strains supports the fact that *PWL3* does not function in preventing the host expanding to grass as *PWL2* does, as it was reported in [70]. On the other hand, the presence of *Avr* genes could be recognized by the host harboring the corresponding *R* genes, thus granting such host with race resistance to the particular strains. In such circumstance, TE insertion to disrupt the *Avr* gene function is a strategy used by the fungal strains to escape such gene-for-gene resistance and shift from avirulent to virulent [85,86].

Overall, we proposed that variations in *Avr* genes profiles may correlate with the host range of *P. oryzae* strains, so that based on analysis of DNA or amino acid sequences of some chosen *Avr* genes, a host range of a particular *P. oryzae* may be predicted. This hypothesis awaits further validation by experiments. Simultaneously, enhancing the monitoring of *Avr* gene variations in field populations of *P. oryzae* will not only deepen our understanding of the molecular mechanisms underlying its host selection and adaptive evolution but also provide a critical decision-making basis for the rational deployment and rotation of resistant rice varieties, thereby safeguarding regional rice production security. With the publication of the genomes of Baicao series [accession numbers: PRJNA1395695], other researchers could also perform further analyses with this information to investigate the molecular mechanisms underlying host adaptation of the *P. oryzae* population.

4. Conclusions

In summary, this study, by integrating phenotypic and genomic data, delineated the differentiation landscape of host adaptation in *P. oryzae* strains infecting barnyard grass and rice. This differentiation resulted from the combined effects of the *Avr* gene repertoire, genomic plasticity (particularly TEs activity), and differences in specific pathogenesis-related gene modules. As a common weed in rice agroecosystems, barnyard grass harbors pathogen populations that constitute a unique genetic pool, for which this study provides genomic resources. Our findings emphasized the importance of focusing on grass hosts and their associated pathogen populations in the integrated management of rice blast.

5. Materials and Methods

5.1. Isolation of Pyricularia Strains

In 2024, leaf samples of barnyard grass exhibiting rice blast symptoms were collected from the Yangjiang Institute of Agricultural Sciences (Baisha) Experimental Base (111°50′ E, 21°52′ N). Single-spore isolation was performed to obtain the *Pyricularia* "Baicao" series of monosporic isolates [87]. Leaf samples containing necrotic lesions were collected using scissors. The leaves were immersed in a 1.2% sodium hypochlorite solution for 3 min, followed by thorough rinsing with sterile water several times until the chlorine odor dissipated. Using sterile forceps, the leaves were transferred to a Petri dish. Unaffected portions of the leaves were excised and discarded, while the remaining segments containing necrotic lesions were placed in a moist chamber. They were then incubated at 25 °C under a 16 h light/8 h dark photoperiod for 3–5 days. During incubation, the leaf segments were observed daily for the emergence of *Pyricularia* mycelia and spores. Under a dissecting microscope, a sterilized fine needle was used to pick a single *Pyricularia* conidium. The picked single spore was transferred onto a RPA medium supplemented with 50 μg/mL streptomycin. The plate was sealed with Parafilm and incubated in darkness at 28 °C for 3–5 days, until a single small colony appeared. Subsequently, mycelia from the edge of the colony were picked with a sterile toothpick and transferred onto a fresh RPA medium containing 50 μg/mL streptomycin for a round of purification. To ensure the purity of the single-spore isolates, *P. oryzae* Strains GDYJ7 and ZJX18, which were previously single-spore isolates from diseased rice plants in Guangdong and Zhejiang provinces, respectively, and preserved in the laboratory, were used as references. Six isolates were cultured on rice polish agar medium (RPA; 20 g/L rice polish, 2 g/L yeast extract, and 12 g/L agar) overlaid with sterile filter papers and cultured in darkness at 28 °C for 14 days until the mycelia fully covered the filter papers. The colonized filter papers were then aseptically collected with sterile forceps, air-dried at 28 °C, and stored at −20 °C [36].

5.2. Pathogenicity Assay

Rice blast fungal isolates were cultured on RPA media at 28 °C with 12 h light followed by 12 h dark treatment for producing mycelium. Seven days later, the mycelia were scraped with a cell spreader and the plates were exposed to continuous light at 28 °C for 3–5 days to promote sporulation. Then, the conidia were washed with 5 mL double-distilled water (ddH_2O) per Petri dish to collect them, and the suspension was adjusted to the concentration of 1×10^5 conidia/mL [88].

The barnyard grass and rice seeds were immersed in water for 2 days at 35 °C in darkness for germination and then grown in the greenhouse with 28–30 °C day/night temperature and 14 h/10 h light/dark period. At three-leaf to five-leaf seedling stages, the second youngest leaves were detached and placed on the ddH_2O with 6-benzylaminopurinehydrochloride (6-BA) in a square Petri dish after punch wounded. Detached leaves of barnyard grass (Echinochloa phyllopogon) or rice cultivars (ZH11, NPB, and CO39) were inoculated with 10 μL of spore suspension. The inoculated leaves were first incubated in darkness at 28 °C and 90% relative humidity for 24 h and then transferred under an alternate dark–light cycle (12 h:12 h) at 28 °C for an additional 4–5 days. Disease symptoms were then recorded and photographed. The experiment included three independent biological replicates [89,90].

5.3. DNA Extraction, Amplification and Sequencing

To prepare the genomic DNA for sequencing, the six isolate strains (Baicao series, GDYJ7, and ZJX18) were cultured in the liquid simple complete medium (CM; 6 g/L casaminoacid, 6 g/L yeast extract, and 10 g/L sucrose) in a 120 rpm shaker at 28 °C for

3–4 days. The mycelia were then collected for the preparation of genomic DNA using the CTAB method [91,92]. The *Avr* sequences of all six *Pyricularia* strains were amplified by PCR using Green Taq Mix (Vazyme, Nanjing, China) and the primers listed in Table 5, with the following amplification conditions: 1 cycle at 95 °C for 3 min, 30 cycles at 95 °C for 15 s, 62 °C for 15 s, and 72 °C for 60 s/kb, followed by 1 cycle at 72 °C for 5 min. PCR amplicons were resolved in 1.5% (*w*/*v*) agarose gel through electrophoresis.

The whole genome was sequenced via massively parallel sequencing (MPS) on the Illumina NovaSeq 6000 platform, by Novogene (Beijing) Co., Ltd. (Beijing, China).

5.4. Assembly, Prediction and Annotation

In this study, de novo whole-genome assembly was performed using SPAdes v4.1.0 on quality-controlled sequencing reads, with assembly parameters set to-k31, 51, 71, 91 [93]. To ensure consistency of the assembly results, experimental replicates were conducted for validation. The completeness of the assembled genomes was assessed with BUSCO v5.8.3 [94]. The average nucleotide identity (ANI) between the sequenced strains and published genomes of closely related species in Table 7 was calculated using FastANI v1.34, allowing phylogenetic relationships to be evaluated at the whole-genome level [95].

Table 7. Strains GenBank whole genome assembly accession number.

Strains	GenBank Accession Number
P. oryzae 131002	GCA_049355375.1
P. oryzae 131021	GCA_049355335.1
P. oryzae 70-15	GCF_000002495.2
P. oryzae Y34	GCA_000292585.1
P. oryzae ZJG1	GCA_025135345.1
P. pennisetigena Br36	GCF_004337985.1
P. grisea D1/s49	GCA_024704135.1
P. grisea Nl907	GCF_004355905.1
P. oryzae D10/s71	GCA_024704055.1
P. oryzae K17	GCA_024704195.1
P. oryzae D15/s47	GCA_024704025.1
P. oryzae K65/159w	GCA_024704275.1
P. oryzae E34	GCA_021845515.1
P. oryzae MZ5-1-6	GCA_004346965.1
P. oryzae D15/s6	GCA_024704165.1
P. oryzae D10/s9	GCA_024704035.1
P. grisea SCAU-2	GCA_040113025.1

Based on the assembled genomes, homology-based gene prediction was performed using protein sequences from closely related species with the Braker v3.0.8 pipeline. Braker is an automated genome annotation pipeline for eukaryotes that integrates tools such as GeneMark and AUGUSTUS [96]. tRNA and rRNA were predicted using tRNAscan-SE V2.0.12 and barrnap v0.9, respectively. For the annotation of non-coding RNAs including snRNA, miRNA, and sRNA, the assembled genomes were compared against the Rfam database (http://rfam.xfam.org/, accessed on 21 October 2025) using Infernal v1.1.5 with its cmsearch program under default parameters [97].

For repetitive sequence prediction, a species-specific repeat library was first constructed using RepeatModeler v2.0.6 to identify repeat types within the library. This library was then merged with the Repbase database (https://www.girinst.org/server/RepBase/index.php, accessed on 7 April 2025) to build the final repeat database. The genome sequences were aligned against this database using RepeatMasker v4.1.8 to predict interspersed repeats [98], while tandem repeats were predicted with TRF v4.09.1 [99].

Protein sequence functional annotation was performed using DIAMOND v2.0.14.152, with alignments conducted against the following databases: KOG (https://www.ncbi.nlm.nih.gov/research/COG/, accessed on 22 May 2025), GO (https://www.geneontology.org/, accessed on 25 May 2025), KEGG (https://www.kegg.jp/, accessed on 7 September 2025), TCDB (http://www.tcdb.org, accessed on 22 May 2025), Pfam (http://pfam.xfam.org/, accessed on 23 May 2025), and CAZy (http://www.cazy.org, accessed on 9 April 2025). The alignment parameter was set to an E-value cutoff of 1×10^{-10}.

Pathogen virulence factors were predicted by aligning protein sequences against the PHI database (http://www.phi-base.org/, accessed on 19 June 2025) and the Database of Fungal Virulence Factors (DFVF) (https://doi.org/10.6084/m9.figshare.31239886, accessed on 1 Februay 2026), with the parameters set to an E-value cutoff of 1×10^{-10} and a sequence identity threshold of >60%.

Secondary metabolite biosynthetic genes and gene clusters were identified using antiSMASH v8.0.0 (https://fungismash.secondarymetabolites.org/, accessed on 10 December 2025) [100], which performs predictions based on profile hidden Markov models (HMMs) specific to biosynthetic types. The presence of core biosynthetic genes in the genomes was further validated by BLAST v2.17.0 analysis, with parameters set to an E-value cutoff of 1×10^{-5} and a sequence coverage threshold of 75%.

5.5. Phylogenetic Analysis and Comparative Genomic Analysis

In this study, we employed OrthoFinder v3.0.1 to perform gene family clustering analysis on the genomes of the six isolated strains [101], along with those of 15 reference strains downloaded from the NCBI genome database (https://www.ncbi.nlm.nih.gov/genome, accessed on 4 December 2025). Orthologous single-copy protein sequences were extracted from the clustering results and aligned using MAFFT v7.525 [102]. A phylogenetic tree was subsequently constructed under the maximum-likelihood (ML) criterion with IQ-TREE v2.4.0, with model selection set to AUTO and branch support assessed via 1000 bootstrap replicates [103].

5.6. Statistical Analysis

Section 2.2 (Pathogenicity Analysis of the Isolated Strains): Lesion area was assessed as described, statistical analyses were performed with GraphPad_Prism 8. DPS 9.01 software, and one-way analysis of variance (ANOVA) tests or Student's *t*-test was carried out. p values < 0.05 were considered as statistically significance.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/jof12020109/s1, Figure S1: Agarose gel electrophoresis image of amplified *Avr* genes from isolated strains; Figure S2: Sequence alignment of *Avr-Pib*; Figure S3: Sequence alignment of *Avr-Pizt*; Figure S4: Sequence alignment of *PWL3*; Table S1: Average nucleotide identity (ANI) analysis; Table S2: The coding genes of the six tested strains with sequence identity higher than 90% to those in the PHI database; Table S3: Comparison of the predicted virulence genes in six tested strains (identity > 90%); Table S4: Distribution of plant cell wall degradation enzymes in six tested strains.

Author Contributions: Y.D., Z.L. and X.Z. (Xiaofan Zhou) conceptualized the study. Y.D. acquired funding. W.S., Z.Z. and X.X. isolated and characterized the *Pyricularia* single-spore strains. W.S., Z.Z., X.X., T.S., S.T. and B.L. helped in conducting experiments and analyzing data. J.W. supplied the *Avr* gene primers. X.Z. (Xiaohan Zhang) and X.Z. (Xiaofan Zhou) conducted the whole-genome sequencing and evolutionary analysis. W.S., X.Z. (Xiaohan Zhang) and Y.D. drafted the original manuscript. W.S. and X.Z. (Xiaohan Zhang) revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: This work is supported by the National Key R&D Program of China (2023YFD1400200). The funders had no role in study design, data collection, and interpretation, or the decision to submit the work for publication.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The all strains whole-genome sequences data have been deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under accession number PRJNA1395695. Supporting data for this study are available in the Supplementary Information.

Acknowledgments: We thank Jiafeng Wang (SCAU) for providing the *Avr* gene primers.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

ANI: average nucleotide identity; NCBI: National Center for Biotechnology Information; RPA: rice polish agar medium; LINEs: long interspersed nuclear elements; SINEs: short interspersed nuclear elements; LTR: long terminal repeat; ncRNAs: non-coding RNAs; tRNAs: transfer RNAs; rRNAs: ribosomal RNAs; snRNAs: small nuclear RNAs; sRNAs: small RNAs; PHI: pathogen–host interactions; DFVF: database of fungal virulence factors; CAZy: carbohydrate-active enzymes; PCWDEs: plant cell wall-degrading enzymes; GHs: glycoside hydrolases; GTs: glycosyltransferases; PLs: polysaccharide lyases; ABC: ATP-binding cassette; MFS: major facilitator superfamily; MDR: multidrug resistance; PDR: pleiotropic drug resistance; ML: maximum-likelihood; TEs: transposable elements; ddH_2O: double-distilled water; 6-BA: 6-benzylaminopurinehydrochloride; MPS: massively parallel sequencing; HMMs: hidden Markov models; FHB: Fusarium head blight; DPs: diterpenoid phytoalexins.

References

1. Fukagawa, N.K.; Ziska, L.H. Rice: Importance for Global Nutrition. *J. Nutr. Sci. Vitaminol.* **2019**, *65*, S2–S3. [CrossRef]
2. Talbot, N.J. Appressoria. *Curr. Biol.* **2019**, *29*, R144–R146. [CrossRef] [PubMed]
3. Foster, A.J.; Talbot, N.J. Getting a grip on blast. *Nat. Microbiol.* **2020**, *5*, 1457–1458. [CrossRef] [PubMed]
4. Wilson, R.A. *Magnaporthe oryzae*. *Trends Microbiol.* **2021**, *29*, 663–664. [CrossRef]
5. Liu, J.; Wang, X.; Mitchell, T.; Hu, Y.; Liu, X.; Dai, L.; Wang, G.L. Recent progress and understanding of the molecular mechanisms of the rice-*Magnaporthe oryzae* interaction. *Mol. Plant Pathol.* **2010**, *11*, 419–427. [CrossRef] [PubMed]
6. Zhang, H.; Wu, Z.; Wang, C.; Li, Y.; Xu, J.R. Germination and infectivity of microconidia in the rice blast fungus *Magnaporthe oryzae*. *Nat. Commun.* **2014**, *5*, 4518. [CrossRef]
7. Dean, R.A.; Talbot, N.J.; Ebbole, D.J.; Farman, M.L.; Mitchell, T.K.; Orbach, M.J.; Thon, M.; Kulkarni, R.; Xu, J.R.; Pan, H.; et al. The genome sequence of the rice blast fungus Magnaporthe grisea. *Nature* **2005**, *434*, 980–986. [CrossRef]
8. Surovy, M.Z.; Islam, T.; von Tiedemann, A. Role of seed infection for the near and far distance dissemination of wheat blast caused by *Magnaporthe oryzae* pathotype Triticum. *Front. Microbiol.* **2023**, *14*, 1040605. [CrossRef]
9. Chung, H.; Goh, J.; Han, S.S.; Roh, J.H.; Kim, Y.; Heu, S.; Shim, H.K.; Jeong, D.G.; Kang, I.J.; Yang, J.W. Comparative Pathogenicity and Host Ranges of *Magnaporthe oryzae* and Related Species. *Plant Pathol. J.* **2020**, *36*, 305–313. [CrossRef]
10. Couch, B.C.; Fudal, I.; Lebrun, M.H.; Tharreau, D.; Valent, B.; van Kim, P.; Notteghem, J.L.; Kohn, L.M. Origins of host-specific populations of the blast pathogen *Magnaporthe oryzae* in crop domestication with subsequent expansion of pandemic clones on rice and weeds of rice. *Genetics* **2005**, *170*, 613–630. [CrossRef]
11. Kellogg, E.A. Evolutionary history of the grasses. *Plant Physiol.* **2001**, *125*, 1198–1205. [CrossRef]
12. Zhong, Z.; Norvienyeku, J.; Chen, M.; Bao, J.; Lin, L.; Chen, L.; Lin, Y.; Wu, X.; Cai, Z.; Zhang, Q.; et al. Directional Selection from Host Plants Is a Major Force Driving Host Specificity in *Magnaporthe* Species. *Sci. Rep.* **2016**, *6*, 25591. [CrossRef] [PubMed]
13. Ganesan, S.; Singh, H.S.; Petikam, S.; Biswal, D. Pathological Status of Pyricularia angulata Causing Blast and Pitting Disease of Banana in Eastern India. *Plant Pathol. J.* **2017**, *33*, 9–20. [CrossRef] [PubMed]
14. Klaubauf, S.; Tharreau, D.; Fournier, E.; Groenewald, J.Z.; Crous, P.W.; de Vries, R.P.; Lebrun, M.H. Resolving the polyphyletic nature of Pyricularia (Pyriculariaceae). *Stud. Mycol.* **2014**, *79*, 85–120. [CrossRef]
15. Stukenbrock, E.H.; McDonald, B.A. The origins of plant pathogens in agro-ecosystems. *Annu. Rev. Phytopathol.* **2008**, *46*, 75–100. [CrossRef]

16. Singh, R.P.; Hodson, D.P.; Huerta-Espino, J.; Jin, Y.; Bhavani, S.; Njau, P.; Herrera-Foessel, S.; Singh, P.K.; Singh, S.; Govindan, V. The emergence of Ug99 races of the stem rust fungus is a threat to world wheat production. *Annu. Rev. Phytopathol.* **2011**, *49*, 465–481. [CrossRef] [PubMed]
17. Corredor-Moreno, P.; Saunders, D.G.O. Expecting the unexpected: Factors influencing the emergence of fungal and oomycete plant pathogens. *New Phytol.* **2020**, *225*, 118–125. [CrossRef]
18. Ceresini, P.C.; Castroagudin, V.L.; Rodrigues, F.A.; Rios, J.A.; Aucique-Perez, C.E.; Moreira, S.I.; Croll, D.; Alves, E.; de Carvalho, G.; Maciel, J.L.N.; et al. Wheat blast: From its origins in South America to its emergence as a global threat. *Mol. Plant Pathol.* **2019**, *20*, 155–172. [CrossRef]
19. Farman, M.; Peterson, G.; Chen, L.; Starnes, J.; Valent, B.; Bachi, P.; Murdock, L.; Hershman, D.; Pedley, K.; Fernandes, J.M.; et al. The Lolium Pathotype of *Magnaporthe oryzae* Recovered from a Single Blasted Wheat Plant in the United States. *Plant Dis.* **2017**, *101*, 684–692. [CrossRef]
20. Barragan, A.C.; Latorre, S.M.; Mock, P.G.; Harant, A.; Win, J.; Malmgren, A.; Burbano, H.A.; Kamoun, S.; Langner, T. Wild grass isolates of Magnaporthe (Syn. Pyricularia) spp. from Germany can cause blast disease on cereal crops. *bioRxiv* **2022**. [CrossRef]
21. Bao, J.; Chen, M.; Zhong, Z.; Tang, W.; Lin, L.; Zhang, X.; Jiang, H.; Zhang, D.; Miao, C.; Tang, H.; et al. PacBio Sequencing Reveals Transposable Elements as a Key Contributor to Genomic Plasticity and Virulence Variation in *Magnaporthe oryzae*. *Mol. Plant* **2017**, *10*, 1465–1468. [CrossRef]
22. Gowda, M.; Shirke, M.D.; Mahesh, H.B.; Chandarana, P.; Rajamani, A.; Chattoo, B.B. Genome analysis of rice-blast fungus *Magnaporthe oryzae* field isolates from southern India. *Genom. Data* **2015**, *5*, 284–291. [CrossRef]
23. Xue, M.; Yang, J.; Li, Z.; Hu, S.; Yao, N.; Dean, R.A.; Zhao, W.; Shen, M.; Zhang, H.; Li, C.; et al. Comparative analysis of the genomes of two field isolates of the rice blast fungus *Magnaporthe oryzae*. *PLoS Genet.* **2012**, *8*, e1002869. [CrossRef] [PubMed]
24. Dong, Y.; Li, Y.; Zhao, M.; Jing, M.; Liu, X.; Liu, M.; Guo, X.; Zhang, X.; Chen, Y.; Liu, Y.; et al. Global genome and transcriptome analyses of *Magnaporthe oryzae* epidemic isolate 98-06 uncover novel effectors and pathogenicity-related genes, revealing gene gain and lose dynamics in genome evolution. *PLoS Pathog.* **2015**, *11*, e1004801. [CrossRef]
25. Chen, C.; Lian, B.; Hu, J.; Zhai, H.; Wang, X.; Venu, R.C.; Liu, E.; Wang, Z.; Chen, M.; Wang, B.; et al. Genome comparison of two *Magnaporthe oryzae* field isolates reveals genome variations and potential virulence effectors. *BMC Genom.* **2013**, *14*, 887. [CrossRef]
26. Yoshida, K.; Saunders, D.G.; Mitsuoka, C.; Natsume, S.; Kosugi, S.; Saitoh, H.; Inoue, Y.; Chuma, I.; Tosa, Y.; Cano, L.M.; et al. Host specialization of the blast fungus *Magnaporthe oryzae* is associated with dynamic gain and loss of genes linked to transposable elements. *BMC Genom.* **2016**, *17*, 370. [CrossRef] [PubMed]
27. Langner, T.; Harant, A.; Gomez-Luciano, L.B.; Shrestha, R.K.; Malmgren, A.; Latorre, S.M.; Burbano, H.A.; Win, J.; Kamoun, S. Genomic rearrangements generate hypervariable mini-chromosomes in host-specific isolates of the blast fungus. *PLoS Genet.* **2021**, *17*, e1009386. [CrossRef]
28. Zheng, H.; Zhong, Z.; Shi, M.; Zhang, L.; Lin, L.; Hong, Y.; Fang, T.; Zhu, Y.; Guo, J.; Zhang, L.; et al. Comparative genomic analysis revealed rapid differentiation in the pathogenicity-related gene repertoires between *Pyricularia oryzae* and Pyricularia penniseti isolated from a Pennisetum grass. *BMC Genom.* **2018**, *19*, 927. [CrossRef]
29. Zhang, N.; Li, X.; Ming, L.; Sun, W.; Xie, X.; Zhi, C.; Zhou, X.; Wen, Y.; Liang, Z.; Deng, Y. Comparative Genomics and Pathogenicity Analysis of Three Fungal Isolates Causing Barnyard Grass Blast. *J. Fungi* **2024**, *10*, 868. [CrossRef] [PubMed]
30. Poppe, S.; Dorsheimer, L.; Happel, P.; Stukenbrock, E.H. Rapidly Evolving Genes Are Key Players in Host Specialization and Virulence of the Fungal Wheat Pathogen Zymoseptoria tritici (Mycosphaerella graminicola). *PLoS Pathog.* **2015**, *11*, e1005055. [CrossRef]
31. Hartmann, F.E.; Sanchez-Vallet, A.; McDonald, B.A.; Croll, D. A fungal wheat pathogen evolved host specialization by extensive chromosomal rearrangements. *ISME J.* **2017**, *11*, 1189–1204. [CrossRef]
32. Lin, L.; Sun, T.; Guo, J.; Lin, L.; Chen, M.; Wang, Z.; Bao, J.; Norvienyeku, J.; Zhang, D.; Han, Y.; et al. Transposable elements impact the population divergence of rice blast fungus *Magnaporthe oryzae*. *mBio* **2024**, *15*, e0008624. [CrossRef]
33. Zhang, N.; Cai, G.; Price, D.C.; Crouch, J.A.; Gladieux, P.; Hillman, B.; Khang, C.H.; LeBrun, M.H.; Lee, Y.H.; Luo, J.; et al. Genome wide analysis of the transition to pathogenic lifestyles in Magnaporthales fungi. *Sci. Rep.* **2018**, *8*, 5862. [CrossRef] [PubMed]
34. Van der Biezen, E.A.; Jones, J.D. Plant disease-resistance proteins and the gene-for-gene concept. *Trends Biochem. Sci.* **1998**, *23*, 454–456. [CrossRef] [PubMed]
35. Younas, M.U.; Wang, G.; Du, H.; Zhang, Y.; Ahmad, I.; Rajput, N.; Li, M.; Feng, Z.; Hu, K.; Khan, N.U.; et al. Approaches to Reduce Rice Blast Disease Using Knowledge from Host Resistance and Pathogen Pathogenicity. *Int. J. Mol. Sci.* **2023**, *24*, 4985. [CrossRef]
36. Hu, Z.J.; Huang, Y.Y.; Lin, X.Y.; Feng, H.; Zhou, S.X.; Xie, Y.; Liu, X.X.; Liu, C.; Zhao, R.M.; Zhao, W.S.; et al. Loss and Natural Variations of Blast Fungal Avirulence Genes Breakdown Rice Resistance Genes in the Sichuan Basin of China. *Front. Plant Sci.* **2022**, *13*, 788876. [CrossRef]

37. Huang, J.; Si, W.; Deng, Q.; Li, P.; Yang, S. Rapid evolution of avirulence genes in rice blast fungus *Magnaporthe oryzae*. *BMC Genet.* **2014**, *15*, 45. [CrossRef] [PubMed]
38. Chuma, I.; Isobe, C.; Hotta, Y.; Ibaragi, K.; Futamata, N.; Kusaba, M.; Yoshida, K.; Terauchi, R.; Fujita, Y.; Nakayashiki, H.; et al. Multiple translocation of the AVR-Pita effector gene among chromosomes of the rice blast fungus *Magnaporthe oryzae* and related species. *PLoS Pathog.* **2011**, *7*, e1002147. [CrossRef]
39. Singh, P.K.; Thakur, S.; Rathour, R.; Variar, M.; Prashanthi, S.K.; Singh, A.K.; Singh, U.D.; Sharma, V.; Singh, N.K.; Sharma, T.R. Transposon-based high sequence diversity in Avr-Pita alleles increases the potential for pathogenicity of *Magnaporthe oryzae* populations. *Funct. Integr. Genom.* **2014**, *14*, 419–429. [CrossRef]
40. Li, W.; Wang, B.; Wu, J.; Lu, G.; Hu, Y.; Zhang, X.; Zhang, Z.; Zhao, Q.; Feng, Q.; Zhang, H.; et al. The *Magnaporthe oryzae* avirulence gene AvrPiz-t encodes a predicted secreted protein that triggers the immunity in rice mediated by the blast resistance gene Piz-t. *Mol. Plant Microbe Interact.* **2009**, *22*, 411–420. [CrossRef]
41. Olukayode, T.; Quime, B.; Shen, Y.C.; Yanoria, M.J.; Zhang, S.; Yang, J.; Zhu, X.; Shen, W.C.; von Tiedemann, A.; Zhou, B. Dynamic Insertion of Pot3 in AvrPib Prevailing in a Field Rice Blast Population in the Philippines Led to the High Virulence Frequency Against the Resistance Gene Pib in Rice. *Phytopathology* **2019**, *109*, 870–877. [CrossRef] [PubMed]
42. Gu, F.; Han, Z.; Zou, X.; Xie, H.; Chen, C.; Huang, C.; Guo, T.; Wang, J.; Wang, H. Unveiling the Role of RNA Recognition Motif Proteins in Orchestrating Nucleotide-Binding Site and Leucine-Rich Repeat Protein Gene Pairs and Chloroplast Immunity Pathways: Insights into Plant Defense Mechanisms. *Int. J. Mol. Sci.* **2024**, *25*, 5557. [CrossRef] [PubMed]
43. Gu, F.; Xie, H.; Huang, Q.; Zhou, W.; Zou, X.; Han, Z.; Guo, T.; Wang, H.; Wang, J. Co-Expression Pattern Analysis of Head-to-Head NLR Gene Pair Pik-H4. *Plant Cell Environ.* **2025**, *48*, 5342–5356. [CrossRef] [PubMed]
44. Biscotti, M.A.; Olmo, E.; Heslop-Harrison, J.S. Repetitive DNA in eukaryotic genomes. *Chromosome Res.* **2015**, *23*, 415–420. [CrossRef]
45. Colonna Romano, N.; Fanti, L. Transposable Elements: Major Players in Shaping Genomic and Evolutionary Patterns. *Cells* **2022**, *11*, 1048. [CrossRef]
46. Metanat, Y.; Sviridova, M.; Al-Nuaimi, B.N.; Janbazi, F.; Jalali, M.; Ghalamkarpour, N.; Khodabandehloo, E.; Ahmadi, E. The role of non-coding RNAs in the regulation of cell death pathways in melanoma. *Discov. Oncol.* **2025**, *16*, 1063. [CrossRef]
47. Hage, H.; Rosso, M.N. Evolution of Fungal Carbohydrate-Active Enzyme Portfolios and Adaptation to Plant Cell-Wall Polymers. *J. Fungi* **2021**, *7*, 185. [CrossRef]
48. Jacob, A.; Willet, A.H.; Igarashi, M.G.; El Hariri El Nokab, M.; Turner, L.A.; Alsanad, A.K.A.; Wang, T.; Gould, K.L. Alpha-glucan remodeling by GH13-domain enzymes shapes fungal cell wall architecture. *Proc. Natl. Acad. Sci. USA* **2025**, *122*, e2505509122. [CrossRef]
49. Wan, Y.; Wang, M.; Chan, E.W.C.; Chen, S. Membrane Transporters of the Major Facilitator Superfamily Are Essential for Long-Term Maintenance of Phenotypic Tolerance to Multiple Antibiotics in *E. coli*. *Microbiol. Spectr.* **2021**, *9*, e0184621. [CrossRef]
50. Doehlemann, G.; Okmen, B.; Zhu, W.; Sharon, A. Plant Pathogenic Fungi. *Microbiol. Spectr.* **2017**, *5*, 701–726. [CrossRef]
51. Mishra, J.; Srivastava, R.; Trivedi, P.K.; Verma, P.C. Effect of virus infection on the secondary metabolite production and phytohormone biosynthesis in plants. *3 Biotech.* **2020**, *10*, 547. [CrossRef]
52. Feng, M.; Yaling, Z.; Xuehui, J.; XiaoYu, Z.; Jun, J. Detection and Analysis of *Magnaporthe oryzae* Avirulence Genes AVR-Pib, AVR-Pik and AvrPiz-t in Heilongjiang Province. *Sci. Agric. Sin.* **2019**, *52*, 4262–4273.
53. Xing, J.; Jia, Y.; Peng, Z.; Shi, Y.; He, Q.; Shu, F.; Zhang, W.; Zhang, Z.; Deng, H. Characterization of Molecular Identity and Pathogenicity of Rice Blast Fungus in Hunan Province of China. *Plant Dis.* **2017**, *101*, 557–561. [CrossRef] [PubMed]
54. Feng, M.; Yaling, Z.; Xuehui, J. Detection and Analysis of *Magnaporthe oryzae* Avirulent Gene AVR-Pita and Its Homologous Genes in Heilongjiang Province. *Chin. J. Rice Sci.* **2020**, *34*, 143–149.
55. Liu, R.; YuHan, Z.H.A.O.; ZhongJu, F.U.; XinYi, G.U.; YanXia, W.A.N.G.; XueHui, J.I.N.; Ying, Y.A.N.G.; WeiHuai, W.U.; Zhang, Y. Distribution and Variation of PWL Gene Family in Rice *Magnaporthe oryzae* from Heilongjiang Province and Hainan Province. *Sci. Agric. Sin.* **2023**, *56*, 264–274.
56. Lee, S.; Kim, C. Chromosome-scale genome assembly of Korean goosegrass (*Eleusine indica*). *Sci. Data* **2025**, *12*, 156. [CrossRef] [PubMed]
57. Jatav, P.K.; Sharma, A.; Dahiya, D.K.; Khan, A.; Agarwal, A.; Kothari, S.L.; Kachhwaha, S. Identification of suitable internal control genes for transcriptional studies in Eleusine coracana under different abiotic stress conditions. *Physiol. Mol. Biol. Plants* **2018**, *24*, 793–807. [CrossRef]
58. Pak, D.; You, M.P.; Lanoiselet, V.; Barbetti, M.J. Management of rice blast (*Pyricularia oryzae*): Implications of alternative hosts. *Eur. J. Plant Pathol.* **2021**, *161*, 343–355. [CrossRef]
59. Gladieux, P.; Condon, B.; Ravel, S.; Soanes, D.; Maciel, J.L.N.; Nhani, A.; Chen, L., Jr.; Terauchi, R.; Lebrun, M.H.; Tharreau, D.; et al. Gene Flow between Divergent Cereal- and Grass-Specific Lineages of the Rice Blast Fungus *Magnaporthe oryzae*. *mBio* **2018**, *9*, e01219-17. [CrossRef]

60. Castroagudin, V.L.; Moreira, S.I.; Pereira, D.A.; Moreira, S.S.; Brunner, P.C.; Maciel, J.L.; Crous, P.W.; McDonald, B.A.; Alves, E.; Ceresini, P.C. Pyricularia graminis-tritici, a new Pyricularia species causing wheat blast. *Persoonia* **2016**, *37*, 199–216. [CrossRef]
61. Seidl, M.F.; Kramer, H.M.; Cook, D.E.; Fiorin, G.L.; van den Berg, G.C.M.; Faino, L.; Thomma, B. Repetitive Elements Contribute to the Diversity and Evolution of Centromeres in the Fungal Genus Verticillium. *mBio* **2020**, *11*, e01714-20. [CrossRef]
62. Hassan, A.H.; Mokhtar, M.M.; El Allali, A. Transposable elements: Multifunctional players in the plant genome. *Front. Plant Sci.* **2023**, *14*, 1330127. [CrossRef] [PubMed]
63. Pereira, D.; Oggenfuss, U.; McDonald, B.A.; Croll, D. Population genomics of transposable element activation in the highly repressive genome of an agricultural pathogen. *Microb. Genom.* **2021**, *7*, 000540. [CrossRef]
64. Badet, T.; Tralamazza, S.M.; Feurtey, A.; Croll, D. Recent reactivation of a pathogenicity-associated transposable element is associated with major chromosomal rearrangements in a fungal wheat pathogen. *Nucleic Acids Res.* **2024**, *52*, 1226–1242. [CrossRef]
65. Xia, C.; Qiu, A.; Wang, M.; Liu, T.; Chen, W.; Chen, X. Current Status and Future Perspectives of Genomics Research in the Rust Fungi. *Int. J. Mol. Sci.* **2022**, *23*, 9629. [CrossRef]
66. Peck, L.D.; Llewellyn, T.; Bennetot, B.; O'Donnell, S.; Nowell, R.W.; Ryan, M.J.; Flood, J.; Rodriguez de la Vega, R.C.; Ropars, J.; Giraud, T.; et al. Horizontal transfers between fungal Fusarium species contributed to successive outbreaks of coffee wilt disease. *PLoS Biol.* **2024**, *22*, e3002480. [CrossRef]
67. Lopez Diaz, C.; Ayhan, D.H.; Rodriguez Lopez, A.; Gomez Gil, L.; Ma, L.J.; Di Pietro, A. Transposons and accessory genes drive adaptation in a clonally evolving fungal pathogen. *Nat. Commun.* **2025**, *16*, 6982. [CrossRef] [PubMed]
68. Le Naour-Vernet, M.; Charriat, F.; Gracy, J.; Cros-Arteil, S.; Ravel, S.; Veillet, F.; Meusnier, I.; Padilla, A.; Kroj, T.; Cesari, S.; et al. Adaptive evolution in virulence effectors of the rice blast fungus *Pyricularia oryzae*. *PLoS Pathog.* **2023**, *19*, e1011294. [CrossRef] [PubMed]
69. Vy, T.T.P.; Inoue, Y.; Asuke, S.; Chuma, I.; Nakayashiki, H.; Tosa, Y. The ACE1 secondary metabolite gene cluster is a pathogenicity factor of wheat blast fungus. *Commun. Biol.* **2024**, *7*, 812. [CrossRef]
70. Kubicek, C.P.; Starr, T.L.; Glass, N.L. Plant cell wall-degrading enzymes and their secretion in plant-pathogenic fungi. *Annu. Rev. Phytopathol.* **2014**, *52*, 427–451. [CrossRef]
71. Chiapello, H.; Mallet, L.; Guerin, C.; Aguileta, G.; Amselem, J.; Kroj, T.; Ortega-Abboud, E.; Lebrun, M.H.; Henrissat, B.; Gendrault, A.; et al. Deciphering Genome Content and Evolutionary Relationships of Isolates from the Fungus *Magnaporthe oryzae* Attacking Different Host Plants. *Genome Biol. Evol.* **2015**, *7*, 2896–2912. [CrossRef] [PubMed]
72. Bulasag, A.S.; Camagna, M.; Kuroyanagi, T.; Ashida, A.; Ito, K.; Tanaka, A.; Sato, I.; Chiba, S.; Ojika, M.; Takemoto, D. Botrytis cinerea tolerates phytoalexins produced by Solanaceae and Fabaceae plants through an efflux transporter BcatrB and metabolizing enzymes. *Front. Plant Sci.* **2023**, *14*, 1177060. [CrossRef]
73. Yamamura, C.; Mizutani, E.; Okada, K.; Nakagawa, H.; Fukushima, S.; Tanaka, A.; Maeda, S.; Kamakura, T.; Yamane, H.; Takatsuji, H.; et al. Diterpenoid phytoalexin factor, a bHLH transcription factor, plays a central role in the biosynthesis of diterpenoid phytoalexins in rice. *Plant J.* **2015**, *84*, 1100–1113. [CrossRef] [PubMed]
74. Macheleidt, J.; Mattern, D.J.; Fischer, J.; Netzker, T.; Weber, J.; Schroeckh, V.; Valiante, V.; Brakhage, A.A. Regulation and Role of Fungal Secondary Metabolites. *Annu. Rev. Genet.* **2016**, *50*, 371–392. [CrossRef]
75. Park, C.H.; Chen, S.; Shirsekar, G.; Zhou, B.; Khang, C.H.; Songkumarn, P.; Afzal, A.J.; Ning, Y.; Wang, R.; Bellizzi, M.; et al. The *Magnaporthe oryzae* effector AvrPiz-t targets the RING E3 ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern-triggered immunity in rice. *Plant Cell* **2012**, *24*, 4748–4762. [CrossRef] [PubMed]
76. Park, C.H.; Shirsekar, G.; Bellizzi, M.; Chen, S.; Songkumarn, P.; Xie, X.; Shi, X.; Ning, Y.; Zhou, B.; Suttiviriya, P.; et al. The E3 Ligase APIP10 Connects the Effector AvrPiz-t to the NLR Receptor Piz-t in Rice. *PLoS Pathog.* **2016**, *12*, e1005529. [CrossRef]
77. Bai, P.; Park, C.H.; Shirsekar, G.; Songkumarn, P.; Bellizzi, M.; Wang, G.L. Role of lysine residues of the *Magnaporthe oryzae* effector AvrPiz-t in effector- and PAMP-triggered immunity. *Mol. Plant Pathol.* **2019**, *20*, 599–608. [CrossRef]
78. Kanzaki, H.; Yoshida, K.; Saitoh, H.; Fujisaki, K.; Hirabuchi, A.; Alaux, L.; Fournier, E.; Tharreau, D.; Terauchi, R. Arms race co-evolution of *Magnaporthe oryzae* AVR-Pik and rice Pik genes driven by their physical interactions. *Plant J.* **2012**, *72*, 894–907. [CrossRef]
79. Guo, J.; Wu, Y.; Huang, J.; Yu, K.; Chen, M.; Han, Y.; Zhong, Z.; Lu, G.; Hong, Y.; Wang, Z.; et al. The *Magnaporthe oryzae* effector Avr-PikD suppresses rice immunity by inhibiting an LSD1-like transcriptional activator. *Crop J.* **2024**, *12*, 482–492. [CrossRef]
80. Wang, Z.; Zhong, G.; Zhang, B.; Xie, Y.; Gan, Y.; Tang, D.; Wang, W. Rice blast pathogen effector AvrPib compromises disease resistance by targeting Raf-like protein kinase OsMAPKKK72 to inhibit MAPK signaling. *J. Integr. Plant Biol.* **2025**, *68*, 486–501. [CrossRef]
81. Liu, Z.; Qiu, J.; Shen, Z.; Wang, C.; Jiang, N.; Shi, H.; Kou, Y. The E3 ubiquitin ligase OsRGLG5 targeted by the *Magnaporthe oryzae* effector AvrPi9 confers basal resistance against rice blast. *Plant Commun.* **2023**, *4*, 100626. [CrossRef]
82. Kang, S.; Sweigard, J.A.; Valent, B. The PWL host specificity gene family in the blast fungus Magnaporthe grisea. *Mol. Plant Microbe Interact.* **1995**, *8*, 939–948. [CrossRef]
83. Zhang, S.; Xu, J.R. Effectors and effector delivery in *Magnaporthe oryzae*. *PLoS Pathog.* **2014**, *10*, e1003826. [CrossRef] [PubMed]

84. Were, V.M.; Yan, X.; Foster, A.J.; Sklenar, J.; Langner, T.; Gentle, A.; Sahu, N.; Bentham, A.; Zdrzalek, R.; Ryder, L.S.; et al. The *Magnaporthe oryzae* effector Pwl2 alters HIPP43 localization to suppress host immunity. *Plant Cell* **2025**, *37*, koaf116. [CrossRef]
85. Brunner, P.C.; McDonald, B.A. Evolutionary analyses of the avirulence effector AvrStb6 in global populations of Zymoseptoria tritici identify candidate amino acids involved in recognition. *Mol. Plant Pathol.* **2018**, *19*, 1836–1846. [CrossRef]
86. Wang, Q.; Li, J.; Lu, L.; He, C.; Li, C. Novel Variation and Evolution of AvrPiz-t of *Magnaporthe oryzae* in Field Isolates. *Front. Genet.* **2020**, *11*, 746. [CrossRef]
87. Fagundes, W.C.; Haueisen, J.; Stukenbrock, E.H. Dissecting the Biology of the Fungal Wheat Pathogen Zymoseptoria tritici: A Laboratory Workflow. *Curr. Protoc. Microbiol.* **2020**, *59*, e128. [CrossRef]
88. Guo, M.; Chen, Y.; Du, Y.; Dong, Y.; Guo, W.; Zhai, S.; Zhang, H.; Dong, S.; Zhang, Z.; Wang, Y.; et al. The bZIP transcription factor MoAP1 mediates the oxidative stress response and is critical for pathogenicity of the rice blast fungus *Magnaporthe oryzae*. *PLoS Pathog.* **2011**, *7*, e1001302. [CrossRef]
89. Chen, X.; Jia, Y.; Wu, B.M. Evaluation of Rice Responses to the Blast Fungus *Magnaporthe oryzae* at Different Growth Stages. *Plant Dis.* **2019**, *103*, 132–136. [CrossRef] [PubMed]
90. Sun, W.; Liu, Q.; Chen, H.; Xie, X.; Zhang, Z.; Zeng, Y.; Zhou, J.; Zhou, X.; Jiang, X.; Liang, Z.; et al. Rice phyllospheric Pantoea spp. suppress blast and bacterial blight diseases. *Environ. Microbiome* **2025**, *20*, 137. [CrossRef] [PubMed]
91. Xu, F.; Liu, X.H.; Zhuang, F.L.; Zhu, J.; Lin, F.C. Analyzing autophagy in *Magnaporthe oryzae*. *Autophagy* **2011**, *7*, 525–530. [CrossRef] [PubMed]
92. Xu, F.; Liu, X.; Wang, J. The complete mitochondrial genome of the rice blast fungus *Pyricularia oryzae* Cavara 1892 strain Guy11 and phylogenetic analysis. *Mitochondrial DNA B Resour.* **2023**, *8*, 1036–1040. [CrossRef]
93. Prjibelski, A.; Antipov, D.; Meleshko, D.; Lapidus, A.; Korobeynikov, A. Using SPAdes De Novo Assembler. *Curr. Protoc. Bioinform.* **2020**, *70*, e102. [CrossRef]
94. Manni, M.; Berkeley, M.R.; Seppey, M.; Simao, F.A.; Zdobnov, E.M. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol. Biol. Evol.* **2021**, *38*, 4647–4654. [CrossRef]
95. Yoon, S.H.; Ha, S.M.; Lim, J.; Kwon, S.; Chun, J. A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie Van. Leeuwenhoek* **2017**, *110*, 1281–1286. [CrossRef]
96. Gabriel, L.; Bruna, T.; Hoff, K.J.; Ebel, M.; Lomsadze, A.; Borodovsky, M.; Stanke, M. BRAKER3: Fully automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP, AUGUSTUS and TSEBRA. *bioRxiv* **2024**. [CrossRef] [PubMed]
97. Nawrocki, E.P.; Eddy, S.R. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* **2013**, *29*, 2933–2935. [CrossRef] [PubMed]
98. Tarailo-Graovac, M.; Chen, N. Using RepeatMasker to identify repetitive elements in genomic sequences. *Curr. Protoc. Bioinform.* **2009**, *Chapter 4*, 4.10.1–4.10.14. [CrossRef]
99. Benson, G. Tandem repeats finder: A program to analyze DNA sequences. *Nucleic Acids Res.* **1999**, *27*, 573–580. [CrossRef]
100. Blin, K.; Shaw, S.; Vader, L.; Szenei, J.; Reitz, Z.L.; Augustijn, H.E.; Cediel-Becerra, J.D.D.; de Crecy-Lagard, V.; Koetsier, R.A.; Williams, S.E.; et al. antiSMASH 8.0: Extended gene cluster detection capabilities and analyses of chemistry, enzymology, and regulation. *Nucleic Acids Res.* **2025**, *53*, W32–W38. [CrossRef]
101. Emms, D.M.; Kelly, S. OrthoFinder: Phylogenetic orthology inference for comparative genomics. *Genome Biol.* **2019**, *20*, 238. [CrossRef] [PubMed]
102. Rozewicki, J.; Li, S.; Amada, K.M.; Standley, D.M.; Katoh, K. MAFFT-DASH: Integrated protein sequence and structural alignment. *Nucleic Acids Res.* **2019**, *47*, W5–W10. [CrossRef] [PubMed]
103. Minh, B.Q.; Schmidt, H.A.; Chernomor, O.; Schrempf, D.; Woodhams, M.D.; von Haeseler, A.; Lanfear, R. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **2020**, *37*, 1530–1534. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.